arXiv:1207.4509v1 [cond-mat.quant-gas] 18 Jul 2012

Dynamics of correlations in a dilute Bose gas following an interaction quench

Stefan S. Natu* and Erich J. Mueller
Laboratory of Atomic and Solid State Physics, Cornell University, Ithaca, New York 14853, USA.

We calculate the dynamics of the one and two body correlation functions in a homogeneous Bose gas at zero temperature following a sudden change in the interaction strength, with and without an underlying lattice. We focus on conceptually simple examples that highlight the features arising separately from interactions and band structure. In the continuum, we show that the Bogoliubov spectrum leads to a diffusive evolution of density correlations for short times, and ballistic at long times. In the lattice we find that the correlation functions develop additional oscillations. Moreover, the lattice dispersion induces an additional velocity scale, and some features instead propagate with that velocity. Finally, we discuss the time-evolution of the contact following a quench. Our predictions, which can be readily tested in experiments, serve as a benchmark for considering the dynamics of more complicated systems.

I. INTRODUCTION

New higher resolution imaging techniques are allowing cold atom experiments to probe spatial correlation functions [1, 2]. Recently focus has turned to the dynamics of these correlations following a sudden change in experimental parameters such as lattice depth or interaction strength [3–7]. In many ways, experiments are ahead of theory, as the very paradigms for thinking of these highly nonequilibrium experiments are just being developed. Here we use the time dependent Bogoliubov approximation to study quenches in the weakly interacting limit. We are thereby able to organize the phenomena, for example separating out interaction effects from lattice effects. Although experiments to date have been in a strong or intermediate coupling regime and cannot be directly modeled by our technique, the existing data appears to be consistent with these organizational principles.

There are a number of fundamental issues which can be probed through the time dependence of correlation functions. For example, experiments on lattice bosons have been related directly to questions of causality [6, 8]. A second interesting direction is to make connections to the fluctuation theorems in classical statistical mechanics [9] which relate equilibrium free energy differences to the work done. This latter physics has been studied in mesoscopic solid state systems by looking at the counting statistics of electrons tunneling through quantum dots [11]. A third direction is using these quenches to help understand how isolated quantum systems approach equilibrium, and determine the timescale for the development of long and short range order [12–18]. Our focus is in a fourth direction, namely on how quenches reveal the structure of the excitation spectrum.

We study a weakly interacting Bose gas, both in free space and in a lattice. We use time dependent Bogoliubov theory to calculate how the correlation functions evolve following a sudden change in the interaction strength. We find that in free space the density correlations spread diffusively at short times, crossing over to ballistic at long times. The speed of propagation is simply the sound speed. In a lattice, new features appear which propagate with a speed determined by the bandwidth. The speed of propagation crosses over to the sound speed at long times. We also look at single particle correlations, for example finding that the time for the condensate fraction to reach equilibrium is simply related to both the initial and final chemical potential.

In addition to this "long wavelength" physics, we find that the short length-scale behavior of the correlation functions has interesting structure. Immediately following the quench, the two-particle correlation function develops a divergence which scales as $1/|\mathbf{r}-\mathbf{r}'|^2$. This structure is due to the singular nature of the two-particle relative wave function at short distances [19]. In equilibrium, this singularity has attracted a lot of attention recently following the work of Tan, who was able to relate the short distance structure of the two-particle correlation function in a two-component Fermi gas to the internal energy via a quantity called the "contact" [20–23]. Here we study the dynamics of the contact (and its generalizations) following a quench.

This paper is organized as follows: In Sec II we describe our system and derive the equations of motion governing the dynamics of the correlation functions in the presence and absence of a lattice. In Sec. III, we calculate the dynamics of single-particle correlation function in the continuum after two types of quenches: a sudden quench from a non-interacting state, and a more general quench between two finite values of the interaction. In Sec. IV we describe the dynamics of the two-particle correlation functions both in the continuum and in a 1D lattice. We limit our discussion to length scales comparable to or greater than the healing length of the condensate. By considering simple instructive cases, we illuminate the role of the excitation spectrum, and compare and contrast the lattice from the continuum. In Sec V., we discuss the short-range physics in the two-particle correlation functions, and relate our results to the contact. In Sec. VI, we summarize our results. Throughout this paper, we limit our discussion to sudden quenches at time $t = 0$ between different initial and final states.

---

*Electronic address: ssn8@cornell.edu

II. MODEL

We start with the Hamiltonian for Bose gas interacting with a contact interaction:

$$\mathcal{H} = \sum_{k} \left(\epsilon_k - \mu\right) a_k^\dagger a_k + \frac{g}{2} \sum_{pqk} a^\dagger_{p+q/2} a^\dagger_{k-q/2} a_k a_p \tag{1}$$

where $g = 4\pi a\hbar^2/m > 0$ where a is the s-wave scattering length and a_k is the bosonic annhilation operator. In free space, the dispersion $\epsilon(k) = \frac{\hbar^2 k^2}{2m}$, while in a lattice, we use $\epsilon(k) = -2J\cos(kd)$, where J is the bandwidth and d is the lattice spacing. Throughout, our free space results are in 3D, while our lattice results are in 1D. These choices of dimensionality are somewhat arbitrary.

The density of condensed atoms is given by $n_0 = \langle a_{k=0}\rangle^2$. We perform the standard Bogoliubov transformation wherein we write the operator $a_{k\neq 0}(t) = u_k(t) b_k + v_k^*(t) b^\dagger_{-k}$ where b_k denotes the bosonic annihilation operator for the non-condensed atoms and has no time dependence. The coherence factors $u_k(t)$ and $v_k(t)$ are complex numbers which satisfy $|u_k(t)|^2 - |v_k(t)|^2 = 1$ for all k, at all times. We always work in the regime where the depletion is small, and the condensate density is roughly equal to the total density.

Ignoring all fluctuations beyond quadratic order in the b_k operators, and neglecting the time dependence of n_0, one obtains the usual Bogoliubov Hamiltonian, which is readily diagonalized to yield the equation of motion for u_k and v_k:

$$\begin{pmatrix} u_k(t) \\ v_k(t) \end{pmatrix} = \left[\cos(E_k t)\hat{I} - i\frac{\sin(E_k t)}{E_k}\begin{pmatrix} \epsilon_k + g_f n_0 & g_f n_0 \\ -g_f n_0 & -(\epsilon_k + g_f n_0) \end{pmatrix}\right]\begin{pmatrix} u_k(0) \\ v_k(0) \end{pmatrix} \tag{2}$$

where $E_k = \sqrt{\epsilon(k)(\epsilon(k) + 2g_f n_0)}$ is the Bogoliubov dispersion where $\epsilon_k = \hbar^2 k^2/2m$. Here $u_k(t=0) = \sqrt{\frac{1}{2}\left(1 + \frac{\epsilon_k + g_i n_0}{E_k}\right)}$ and $v_k(t=0) = -\sqrt{\frac{1}{2}\left(\frac{\epsilon_k + g_i n_0}{E_k} - 1\right)}$ are the static solutions. Here g_i and g_f denote the initial and final interactions strengths respectively.

A. Expressions for Correlation functions

Here we are interested in the dynamics of two correlation functions: the non-condensed fraction ($n_{ex} = \sum_{\mathbf{k}\neq\mathbf{0}} \langle a_k^\dagger a_k\rangle$), and the density-density correlation function $g^{(2)}(\mathbf{r} - \mathbf{r}') = \sum_{\mathbf{q}} e^{i\mathbf{q}\cdot(\mathbf{r}-\mathbf{r}')}\langle\rho_{\mathbf{q}}\rho_{-\mathbf{q}}\rangle$ where $\rho_{\mathbf{q}} = \sum_{\mathbf{k}} a^\dagger_{\mathbf{k}+\mathbf{q}} a_{\mathbf{k}}$. The former is readily measured in time-of-flight experiments [7], while the latter is probed using Bragg spectroscopy [4, 21, 24], noise correlations [26] or by direct in-situ measurements [5, 6]. At zero temperature,

$$n_{ex}(t) = \sum_{\mathbf{k}} |v_k(t)|^2 \tag{3}$$

$$g^{(2)}_\delta(t) = n_0^2 + 2n_0 n_{ex}(t) + n_0 \sum_{\mathbf{k}} e^{i\mathbf{k}\cdot\delta}\Big(2|v_k(t)|^2 + u_k^*(t) v_k(t) + u_k(t) v_k^*(t)\Big) + \tilde{g}^{(2)}_\delta \tag{4}$$

where $\delta = |\mathbf{r} - \mathbf{r}'|$.

The term $\tilde{g}^{(2)}_\delta$ is quartic in the u_ks and v_ks and arises from correlations between the non-condensed atoms. These correlations become important only at extremely short distances $\delta \sim a \sim 50$nm, which is roughly 20 times smaller than the typical condensate healing length $\zeta \sim \mu$m. As we are primarily concerned with $\delta \geq \zeta$, we ignore corrections to the dynamics arising from this term, except in Section V.

The first term in Eq. 4 is roughly equal to n^2 where $n = n_0 + n_{ex}$ is the total density. The second term denotes correlations between the condensed and non-condensed atoms. In a non-interacting gas at zero temperature, $u_k = 1$ and $v_k = 0$, therefore $n_{ex} = 0$ and $g^{(2)}_\delta = n_0^2$, independent of δ.

B. Self-consistency

In a more sophisticated model, we would let n_0 be time-dependent. Such an elaboration is unnecessary for quenches where the change in the equilibrium condensate fraction is small. Mathematically the resulting nonlinear theory is much more unwieldy.

III. NON-CONDENSED FRACTION

A. Quench from $g_i = 0$ to $g_f > 0$

We first consider the dynamics of the non-condensed fraction. The conceptually simplest case is to consider a quench from $g = 0$ to some final $g > 0$. From Eq. 2, we find that $v_k(t) = -ign_0 \sin(E_k t)/E_k$, thus $n_{ex} = \sum_{\mathbf{k}} |v_k|^2 = (gn_0)^2 \sum_{\mathbf{k}} \sin(E_k t)^2/E_k^2$. At long times, the sum is dominated by small values of k, where the dispersion is linear $E_k = ck$, where $c = \sqrt{gn_0/m}$ is the speed of sound. With this substitution, the sum is readily performed to yield the expression:

$$n_{ex}(t) = \frac{1}{4\pi}\zeta^{-3}(1 - e^{-4t/\tau_{MF}}) \tag{5}$$

where $\zeta = \hbar/\sqrt{mgn_0}$ is the healing length of the condensate and $\tau_{MF} = \hbar/gn_0$ is the characteristic relaxation time following the quench. Therefore the population of non-condensed particles grows linearly at short times, saturating on timescales $t \sim \tau_{MF}$. At long times, the number of excitations created is *larger* than the predicted zero temperature equilibrium value $n_{ex,eq} = \zeta^{-3}/3\pi^2$. Given that the Bogoliubov approximation is collisionless, there is no reason to expect this distribution to be thermal.

B. Quench from $g_i > 0$ to $g_f > 0$

Next, we consider a quench from some initial interaction strength g_i to a final value g_f. The non-condensed fraction is then

$$n_{ex}(t) = \sum_k |v_k(0)|^2 - g_f(g_i - g_f)n_0^2 \frac{\epsilon(k)\sin(E_k t)^2}{E_k^3} \tag{6}$$

The first term in this expression is the number of excitations present in the system initially. The minus sign in front of the second term indicates that if $g_i > g_f$, the number of non-condensed particles decreases over time, and vice-versa if $g_i < g_f$. Alternatively, the condensate fraction grows if $g_i < g_f$ and reduces if $g_f > g_i$.

The number of *new* excitations created/destroyed is thus proportional to $g_f \times (g_i - g_f)$. If $g_f = 0$, then this term is identically zero and no more excitations are created or lost. The initially quantum depleted superfluid remains depleted, unless interactions are present to redistribute quasi-momentum.

To obtain the relevant timescale governing the redistribution of quasi-momentum, we evaluate the expression for n_{ex} assuming a linear dispersion. Defining $\zeta = \hbar/\sqrt{mg_i n_0}$ as the healing length of the initial state, we obtain:

$$n_{ex}(t) = \frac{1}{3\pi^2}\zeta^{-3}\Big(1 - 3\pi\tilde{g}_f(1-\tilde{g}_f)t/\tau_1\Big[I_0(4t/\tau_2) - L_0(4t/\tau_2)\Big]\Big) \tag{7}$$



FIG. 1: **Quasi-momentum redistribution in 3D following a sudden change in interactions:** Plotted is the fraction of non-condensed atoms normalized to the initial excitation fraction following a sudden quench to weaker interactions $g_f/g_i = 0.6$. The excitation density is expressed in terms of ζ, the healing length of the initial condensate. Note that the analytic formula Eq. 7 (solid curve) captures the dynamics very well, particularly the long time limit. The characteristic relaxation time is set by $\tau_2 = \hbar/g_f n$, the mean-field time of the final state. The dashed curve is the excitation fraction at the final interaction strength in equilibrium.

where $\tilde{g}_f = g_f/g_i$, $\tau_1 = \hbar/\sqrt{g_f g_i} n_0$ and $\tau_2 = \hbar/g_f n_0$, and I_0 and L_0 denote the Bessel function of the second kind and the modified Struve function respectively. For small arguments, $I_0(x) - L_0(x) \approx 1 - 2x/\pi + \mathcal{O}(x^2)$, while for $x \gg 1$, $I_0(x) - L_0(x) \approx 2/\pi x + \mathcal{O}(x^{-3})$.

Unlike the case of a quench from zero interactions, where τ_{MF} was the only relevant timescale, the dynamics here are governed by *two* timescales: at short times, the number of excitations decreases linearly in time with a timescale given by the geometric mean of the initial and final interaction strengths $\tau_1 \propto 1/\sqrt{g_i g_f} n_0$. At long times $t > \tau_2$, the number of excitations created/destroyed saturates to a constant proportional to $\tilde{g}_f(1 - \tilde{g}_f)$.

In Fig. 1 we plot Eq. 7. Also shown is the result of numerically integrating Eq. 6 assuming the full Bogoliubov dispersion, showing that Eq. 7 indeed captures all the relevant features seen in the dynamics.

Again the number of excitations present at long times after the quench is always greater than the number of excitations in the equilibrium final state. This is consistent with a gain in entropy. Note that in the weakly interacting regime discussed here, the fraction of excited particles is typically smaller than the condensate fraction by a factor of $\sqrt{n_0 a^3} \ll 1$, so the deviation of the excited fraction from its equilibrium value may be somewhat difficult to measure experimentally. The clearest (albeit somewhat trivial) demonstration of a non-equilibrium final state is a sudden quench to $g = 0$, where the excitation fraction remains constant at all times following the quench.

IV. TWO-PARTICLE CORRELATIONS

A. Diffusive to ballistic crossover

We now consider the evolution of $g_\delta^{(2)}$ following a sudden quench from $g = 0$ to $g = g_f > 0$. At $t = 0$, $u_k = 1$ and $v_k = 0$, and neglecting the quartic term $\tilde{g}_\delta^{(2)}$, Eq. 4 simplifies to $g_\delta^{(2)}(t) = n^2 - g n_0^2 \sum_{\mathbf{k}} e^{i\mathbf{k}\cdot\delta} \frac{\sin^2(E_k t)}{E_k^2} \epsilon_k$. At long times, the phase of the $\sin(E_k t)^2$ oscillates rapidly and we can simply replace $\sin(E_k t)^2 \to \langle \sin(E_k t)^2 \rangle = 1/2$. We find that $g_\delta^{(2)}(t \to \infty) = n^2 - \frac{n_0 \zeta^{-3}}{4\pi\tilde{\delta}} e^{-2\tilde{\delta}}$, where $\tilde{\delta} = \delta/\zeta$. At long times, $g_\delta^{(2)}$ appears to diverge as $\delta \to 0$ as $1/\delta$. On distances much shorter than ζ, one must also include the contribution from $\tilde{g}_\delta^{(2)}$, which scales as $1/\delta^2$. Therefore, at small distances, $g_\delta^{(2)} \sim 1/\delta^2 - a/\delta$ [19].

Focussing on the longer range physics, we introduce dimensionless variables $\tilde{\delta} = \delta/\zeta$ and $\tilde{t} = t/\tau_{MF}$ and write

$$g_\delta^{(2)} - n^2 = -\frac{1}{\pi^2 \tilde{\delta}} n_0 \zeta^{-3} \int_0^\infty d\tilde{k} \tilde{k} \frac{\sin(\sqrt{2}\tilde{k}\delta)}{\tilde{k}^2 + 2} \times \sin^2\left(\sqrt{\tilde{k}^2(\tilde{k}^2+2)}\tilde{t}\right) \tag{8}$$

where $k = \sqrt{2}\tilde{k}/\zeta$, and we have neglected $\tilde{g}^{(2)}$.

In Fig. 2 we plot the evolution of the density-density correlation function $g^{(2)} - n_0^2$ as a function of $\tilde{t}$ for $\tilde{\delta} = 4$, normalized to its asymptotic value. At short times, the correlations rapidly oscillate. The temporal period of oscillations increases with time. The structure is revealed in a saddle point approximation (with the dominant wave-vectors near $k \sim \delta/t$). Within this approximation, $g_\delta^{(2)}$ oscillates as $\sin(\delta^2/t + \phi)$. At very long times, only particles with $k \approx 0$ contribute to the dynamics and the correlations saturate. As in the case of the one-body correlations, the requirement for correlations to saturate is $(t - \tilde{\delta}/c) \gg \tau_{MF}$.

The temporal location of the last minimum in the correlation function (t_{min}) is denoted by an arrow in Fig. 2(left). On the right, we show how this minimum disperses with $\tilde{\delta}$. When $\delta \leq \zeta = \hbar/\sqrt{m g_f n_0}$, the correlations spread diffusively, while for $\delta \geq \zeta$, correlations spread ballistically. This crossover arises from the nature of the Bogoliubov dispersion. At short distances, the large momentum structure of the dispersion dominates $E_k \sim k^2$, leading to diffusive dynamics; longer range correlations are governed by the linear part of the dispersion and spread ballistically.

To further illustrate the existence of this crossover, we also consider a quench in the reverse direction from $g_i > 0$ to $g_f = 0$. We find that $g_\delta^{(2)}(t) = g^{(2)}(t = 0) - 2n_0 \sum_k e^{i\mathbf{k}\cdot\delta} \sin(\epsilon_k t)^2 u_k(0) v_k(0)$. Note that even for a quench to the non-interacting state, density-density correlations evolve in time (though the condensate fraction remains fixed). As shown in Fig. 2(right), both short and long range correlations spread *diffusively* (we find $\tilde{\delta} = \pi\sqrt{\tilde{t}_{min}}$) as the non-interacting dispersion is $E_k = k^2/2m$ for all k.



FIG. 2: **Dynamics of density-density correlations in 3D** Left: density-density correlations $g^{(2)}(\delta/\zeta = 4) - n^2$ normalized to the asymptotic value at long times for a quench from a non-interacting state to some final $g_f > 0$. Length and time is measured in terms of the condensate healing length ($\zeta = \hbar/\sqrt{m g_f n_0}$) and mean-field time $\tau_{MF} = \hbar/(g_f n_0)$ in the final state. Correlations develop in an oscillatory manner and rapidly saturate at times $t - \delta/c > \tau_{MF}$. Inset shows the temporal location of the minimum in the correlation function (arrow on left graph) (abscissa) plotted versus δ/ζ (ordinate). Long range correlations $\delta/\zeta \gg 1$ spread ballistically while short range correlations $\delta/\zeta \ll 1$ spread diffusively. The lines show purely linear and linear plus square root fit to the data. The latter captures the data accurately. Right: Dots are the same as the inset on left. Solid line shows the location of the minima in the correlation function for a quench from finite interactions to $g = 0$. The dynamics are purely diffusive in this case, with a diffusion constant $\approx \pi$.

Just as with the single particle correlations, for a sudden quench from g_i to g_f, density-density correlations typically *do not* attain their equilibrium values at long times zero temperature values. For example, for a quench from g_i to $g_f = 0$, we find that while at long distances $g_{\delta \gg \zeta}^{(2)}(t = \infty) \to n_0^2 + 2n_0 n_{ex}$ whereas for short distances $g_{\delta \ll \zeta}^{(2)}(t = \infty) \to n_0^2 + 4n_0 n_{ex}$. Interestingly the density-density correlations at long times following a quench to a non-interacting final state do not exhibit the $1/\delta$ divergence at small distances. In Fig. 3 we plot the density-density correlations at $t \to \infty$ after a quench to $g_f = 0$.

B. Lattice vs. continuum

We now ask whether additional features are found in the presence of an underlying lattice. The conceptually simplest case to consider is a quench from finite interactions to $g = 0$, in the presence and absence of a lattice. Here we consider a 1D lattice with band structure $\epsilon(k) = 4J\sin^2(kd/2)$, but the physics described here is qualitatively similar in higher dimensions.

From Eqs. 2 and 4, we find that



FIG. 3: **Long-time density-density correlation function** Density-density correlation function at long times following a quench from finite interaction strength to $g = 0$. The equilibrium value of $g^2(\delta)$ for a non-interacting condensate is n_0^2. Instead at long distances $g^{(2)}_{\delta\gg\zeta}(t=\infty) \to n_0^2 + 2n_0 n_{ex}$ whereas for short distances $g^{(2)}_{\delta\ll\zeta}(t=\infty) \to n_0^2 + 4n_0 n_{ex}$. Correlations fall off exponentially on a length scale proportional to the coherence length of the initial interacting gas.

$$g_\delta^{(2)}(t) = g_\delta^{(2)}(0) - \frac{g_i n_0^2}{2\pi d}\int_{-\pi}^{\pi} dk e^{ik\tilde{\delta}} \frac{\sin(4J\sin^2(k/2)t)^2}{\sin(k/2)\sqrt{4J\sin^2(k/2)+2g_i n_0}} \tag{9}$$

where $\tilde{\delta} = \delta/d$. Normalizing the energy and time units by J, we numerically integrate Eq. 9 and plot in Fig. 4 the typical post-quench dynamics of the density-density correlations as a function of time, following such a quench.

At long times, the correlations in the lattice and continuum both decay to their asymptotic values algebraically (with different exponents), but in addition, the lattice introduces periodic oscillations. The period of these oscillations is largely independent of δ, and is proportional to the time it takes for particles to travel a lattice unit. These oscillations are a purely lattice effect and were also observed experimentally for quenches within the Mott insulating phase [6].

The differences between the lattice and continuum are even more striking at short times, as these reveal the high momentum structure of the underlying excitation spectrum. In Fig. 4 we plot the density-density correlation function for different values of δ and highlight two features. For comparison we plot the density-density correlations in the continuum for the same quench. The open circles denote the position of the first maximum in the correlation function while the filled circles denote the minimum in the correlation function. On the right we show that these features disperse differently, the position of the maximum disperses ballistically, while the position of the minimum disperses diffusively.

The linearly dispersing maximum has no analog in the continuum and emerges purely due to the lattice bandstructure, which imposes a maximum velocity on the growth of correlations. In 1D, this velocity is $4Jd$ [25].

For quenches to an interacting final state $g_f \neq 0$, the dynamics become more complicated. The minimum in the correlation functions disperses diffusively, then ballistically, similar to the 3D continuum case. The maximum in the correlation function disperses linearly with a velocity set by the band structure for small values of g_f crossing over to the superfluid velocity for larger g_f.



FIG. 4: **Dynamics of density-density correlations in a 1D lattice** (Left): Typical structure of density-density correlations $g^{(2)}(\delta)(t) - g^{(2)}(\delta)(0)$ for different values of δ normalized to the asymptotic value at long times for a quench from $g_i > 0$ to $g_f = 0$. Lengths and times are measured in terms of the lattice spacing and inverse hopping J^{-1}. Correlations in a lattice oscillate with a period proportional to $1/J$ and decay with a time constant independent of δ. The temporal location of the first maximum (t_{max}) and the minimum (t_{min}) are indicated by open and filled circles. (Right, Top): A plot of the evolution of the density-density correlations in the 3D continuum for the same quench at $\delta = 5\zeta$ for comparison. The minimum in the correlation function (filled circle) disperses diffusively (Right, Bottom): t_{max} disperses ballistically at all distances, while t_{min} disperses diffusively. This latter features is the analog of the quantity highlighted on the top right.

V. SHORT DISTANCE STRUCTURE OF TWO-PARTICLE CORRELATIONS

Thus far we have neglected the contribution to the correlation functions arising purely from the non-condensed atoms. For $\delta \sim \zeta$, and weak interactions, these terms are of order $\mathcal{O}(a/\zeta)^2$, (for typical densities $a \sim 50$nm and $\zeta \sim \mu$m) and can be ignored. However for $\delta \ll \zeta$, these terms give rise to a $1/\delta^2$ divergence in the correlation function as a result of the singular nature of the relative wave-function of two particles interacting with a zero range interaction [19]. The two-particle correlation function $g^{(2)}(\delta) \sim (1 - a/\delta)^2$ where the correlations between non-condensed particles dominate the short distance behavior and the interaction between the condensed and non-condensed particles leads to a sub leading $1/\delta$ correction which is opposite in sign.

Recently Shina Tan showed that for a two-component



FIG. 5: **Dynamics of the contact $\mathcal{C}(\delta)$ in a 3D Bose gas:** (Color Online) (Left) Dynamics of the position dependent contact $\mathcal{C}(\delta) = \delta^2 g^{(2)}(\delta)$ (normalized to the true contact within the Bogoliubov approximation) at different times following the quench, as a function of δ. At $t < 0$, the gas is non-interacting and $\mathcal{C}(\delta) = 0$. Immediately after the quench, the zero distance correlations respond instantaneously. Colors correspond to different times: $t/\tau_{MF} = 0.025$ (green), 0.1 (red), 0.3 (blue), 0.9 (black). We choose $a = \zeta$. For times $\tau > \tau_{MF}$, $\mathcal{C}(\delta)$ saturates to it's equilibrium value for a given δ. (Right) Dependence of $\mathcal{C}(\delta)$ with scattering length a measured in units of ζ. From top to bottom: $\delta/\zeta = 1$, $\delta/\zeta = 0.1$ and $\delta/\zeta = 0.01$. In order to obtain the expected quadratic dependence of the true contact $\mathcal{C}$ with a for a Bose-Einstein condensate, one has to probe correlations on length scales $\delta \sim$ nm.

Fermi gas interacting with a contact interaction, the short distance structure of the two-body correlation is related to the internal energy via the "contact" defined as $\mathcal{C} = \lim_{\delta\to 0} \delta^2 g^{(2)}(\delta)$ [20]. For a Bose-Einstein condensate, $\mathcal{C}$ to leading order is $16\pi^2 n_0^2 a^2 (1 + \mathcal{O}(\sqrt{na^3}))$ [19, 23]. Experiments using Bragg spectroscopy to probe the Fourier transform of $g^{(2)}$ have a typical momentum resolution of $k = 2\pi/\delta_{min}$ where $\delta_{min} \sim 0.5\mu$m [21]. As a result, experiments actually probe $\mathcal{C}(\delta) = \delta^2 g^{(2)}(\delta)$ evaluated at δ_{min}.

The dominant contribution to these correlations from the non-condensed atoms is a term of the form $\tilde{g}^{(2)}_{\delta} = |\sum_{\mathbf{p}} e^{i\mathbf{p}\cdot\delta} u_p^*(t) v_p(t)|^2$. Adding this term to Eq. 4, and multiplying by δ^2 we numerically integrate the resulting equation to obtain $\mathcal{C}(\delta)$ for a quench from $g_i = 0$ to $g_f = g$.

In Fig. 5 we plot $\mathcal{C}(\delta)$ as a function of δ at different times. The contribution to $g^{(2)}$ from correlations between the condensate and the non-condensate atoms dominates unless $\delta/\zeta \ll 1$ or $a \geq \zeta$. We choose $a = 5\zeta$. Immediately following the quench, the zero range correlations jump to their equilibrium value. However $\mathcal{C}(\delta)$ changes in time for finite δ and relaxes to a stationary value (denoted by the black curve) on times $t \sim \tau_{MF}$. Therefore, although the true contact does not have any time-dependence, the experimentally relevant quantity $\mathcal{C}(\delta)$ changes in time following an interaction ramp.

Finally we remark that *even in equilibrium* the scattering length dependence of $C(\delta)$ is strongly influenced by δ. For example, the $\delta \sim \mu$m scale studied in experiments, one has to go to very large interactions in order to suppress the contribution to $g^{(2)}$ arising from correlations between the condensed and non-condensed atoms. We illustrate this in Fig. 5(right) where we plot $\mathcal{C}(\delta, a)$ as a function of a for different values of δ, after the system has equilibrated following the ramp. We find that for $\delta \sim \zeta$ (typically μm), $\mathcal{C}(a)$ varies linearly with a at small a. The quadratic dependence of $\mathcal{C}(\delta)$ with a for a weakly interacting BEC emerges only for small $\delta \sim$nm.

VI. CONCLUSIONS

Relating the information contained in the fluctuations of an interacting system driven out of equilibrium to the underlying many body parameters is a challenging task. Here we study a simple model, a homogenous Bose gas at zero temperature, where one can calculate the one and two body correlations of the system following a sudden change in the interactions. Our study reveals that the dynamics of these correlations reveal a wealth of information about the underlying excitation spectrum of the system.

We first consider the dynamics of single particle correlations, such as the condensate fraction or equivalently, the excited fraction of atoms following a sudden interaction ramp. For a quench from a non-interacting gas, we find that all the dynamics are governed by a single timescale, the mean-field time. For quenches between two interacting systems, we find two timescales governing the long and short term dynamics of the gas. Moreover we find that the excited fraction at long times after the quench is always greater than the expected equilibrium value.

We then consider the dynamics of the two particle correlation function. For quenches between interacting initial and final states, we find a non-trivial crossover between diffusive spreading of short-range correlations and the ballistic spread of long range correlations. We relate this crossover to the underlying Bogoliubov dispersion. We then consider the case of a Bose gas in a lattice and discuss the additional features that arise in the correlations from the band structure. Notably, we find that the band structure leads to an additional linearly dispersing feature in the correlation functions.

Finally we discuss the dynamics of the contact following a sudden quench. We find that while the true zero range contact instantaneously takes on the new equilibrium value, the finite resolution of an experiment will make the contact appear time dependent.

Acknowledgements.— This work was supported by a grant from the Army Research Office with funding from the DARPA OLE program and by the National Science Foundation through grant No. 1066293.

[1] W. S. Bakr *et al.* Science **329** 547 (2010).
[2] C-L. Hung *et al.* New. J. Phys. **13** 075019 (2011).
[3] M. Greiner *et al* Nature **415** 39 (2002).
[4] H. Miyake, *et al.* Phys. Rev. Lett. **107** 175302 (2011).
[5] S. Trotzky *et al.* Nature Physics **8** 325 (2012).
[6] M. Cheneau *et al.* Nature **481** 484 (2012).
[7] S. S. Natu, D. C. McKay, B. DeMarco and E. J. Mueller, Phys. Rev. A **85** .061601 (R) (2012).
[8] E. H. Lieb and D. W. Robinson, Commun. Math. Phys., **28** 251 (1972).
[9] C. Jarzynski, Phys. Rev. Lett. **78** 2690 (1997); E. Sevick, R. Prabhakar, S. R. Williams and D. J. Searles, Annu. Rev. Phys. Chem **59** 603 (2008).
[10] *Quantum Noise in Mesoscopic Physics* edited by Yu. V. Nazarov (Kluwer, Dordrecht, 2003); B. Reulet, J. Senzier and D. E. Prober, Phys. Rev. Lett. **91** 196601 (2003).
[11] C. Flindt *et al.* Proc. Natl. Acad. Sci. **106** 10116 (2009)
[12] M Gring *et al.* arXiv:1112.0013.
[13] S. S. Natu and E. J. Mueller arXiv: 1201.6674.
[14] A. Polkovnikov, S. Sachdev, S. M. Girvin, Phys. Rev. A **66** 053607 (2002).
[15] M. Rigol and A. Muramatsu, Phys. Rev. Lett. **93** 230404 (2004).
[16] F. M. Cucchietti, B. Damski, J. Dziarmaga and W. Zurek, Phys. Rev. A **75** 023603 (2007).
[17] T. W. B. Kibble, J. Phys. A **9** 1387 (1976); W. H. Zurek, Nature **317** 505 (1985).
[18] U. R. Fischer, R. Schutzhold and M. Uhlmann, Phys. Rev. A **77** 043615 (2008).
[19] T. D. Lee, K. Huang and C. N. Yang, Phys. Rev. **106** 1135 (1957); M. Naraschewski and R. J. Glauber, Phys. Rev. A **59** 4595 (1999).
[20] S. Tan, Ann. Phys. **323** 2952 (2008); 2971 (2008); 2987 (2008); E. Braaten and L. Platter, Phys. Rev. Lett. **100** 205301 (2008); F. Werner, L. Tarruel and Y. Castin Eur. Phys. J B **68** 401 (2009); S. Zhang and A. J Leggett, Phys. Rev. A **79** 023601 (2009).
[21] E. D. Kuhnle *et al.* Phys. Rev. Lett. **105** 070402 (2010).
[22] J. M. Diederix, T. C. F van Heijst and H. T. C Stoof, Phys. Rev A **84** 033618 (2011).
[23] R. J. Wild, *et al.* Phys. Rev. Lett. **108** 145305 (2012).
[24] T. A. Corcovilos, *et al.* Phys. Rev. A **81** 013415 (2010).
[25] P. Calabrese and J. Cardy, Phys. Rev. Lett. **96** 136801 (2006).
[26] E. Altman, E. Demler and M. D. Lukin, Phys. Rev A **70** 013603 (2004); M. Greiner, C. A. Regal J. T. Stewart and D. S. Jin Phys. Rev. Lett. **94** 110401 (2005).